

WEIGHTED ESTIMATES FOR CONIC FOURIER MULTIPLIERS

ANTONIO CÓRDOBA AND KEITH M. ROGERS

ABSTRACT. We prove a weighted inequality which controls conic Fourier multiplier operators in terms of lacunary directional maximal operators. By bounding the maximal operators, this enables us to conclude that the multiplier operators are bounded on $L^p(\mathbb{R}^3)$ with $1 < p < \infty$.

1. INTRODUCTION

For directions $\omega \in \Omega$ contained in the unit circle $\mathbb{S}^1 \subset \mathbb{R}^2$ we form the associated polygon $\mathcal{P}_\Omega$, with sides contained in $\omega + \omega^\perp$ (see Figure 1). We consider the associated cones

$$\Gamma_\Omega = \left\{ \xi \in \mathbb{R}^3 : \frac{(\xi_1, \xi_2)}{|\xi_3|} \in \mathcal{P}_\Omega \right\}$$

and the Fourier multiplier operators T_Ω defined, initially on Schwartz functions, by

$$T_\Omega : f \mapsto \left(\chi_{\Gamma_\Omega} \hat{f} \right)^\vee.$$

Here, $\hat{\cdot}$ and $\vee$ denote the Fourier transform and inverse transform, respectively. The purpose of this note is to provide a condition on the directions Ω which ensures T_Ω is bounded from $L^p(\mathbb{R}^3)$ to $L^p(\mathbb{R}^3)$, where $1 < p < \infty$. The question is only interesting when $p \neq 2$, as when $p = 2$ the operator is bounded by Plancherel's theorem, and no restrictions on the directions are necessary. As shown by Fefferman [15], one cannot take all the directions of the circle, converting the polygon in a disc, when $p \neq 2$, however one can consider the cone associated to the disc if the multiplier is smoothed out appropriately (see [17] and the references therein).

We will control T_Ω , via a weighted inequality, by a combination of directional maximal operators (see the forthcoming Theorem 3.3). This point of view has its origins in a conjecture of the first author [10] for the Bochner–Riesz multiplier (see also [23, 11, 5, 7, 8, 16] for progress on this conjecture

Key words and phrases. Fourier multipliers; maximal operators; lacunary directions.

Supported by the Spanish grants MTM2010-16518, MTM2011-22851, SEV-2011-0087 and the ERC grant 277778.

2010 Mathematics Subject Classification. Primary 42B25; Secondary 26B05.




FIGURE 1. The polygon $\mathcal{P}_\Omega$ and part of the cone Γ_Ω associated to a set Ω of lacunary directions.

and [13, 6, 3, 4] for closely related problems). When the directions are lacunary of finite order (loosely speaking, a lacunary set of order K is a lacunary sequence which has lacunary sets of order $K - 1$ accumulating at each direction; see Definition 3.1) we are able to bound the relevant maximal operators, which yields the following theorem.

Theorem 1.1. *Let $1 < p < \infty$ and suppose that $\Omega \subset \mathbb{S}^1$ is lacunary of finite order. Then T_Ω is bounded from $L^p(\mathbb{R}^3)$ to $L^p(\mathbb{R}^3)$.*

The proof of this can be found in the final section. In the following section, we will recall a weighted inequality for singular integrals, and prove a weighted, angular Littlewood–Paley inequality. That angular square functions are bounded in L^p is well-known (see [13, 18] for lacunary directions and [21] for lacunary directions of finite order), however we will require a weighted version in order to control T_Ω in terms of maximal operators.

2. AN ANGULAR LITTLEWOOD–PALEY INEQUALITY

Theorem 1.1 will be proved by a number of applications of the following one-dimensional result for singular integrals. There are multidimensional versions of some of the results in this section, however we focus only on what we will need. Fundamental to our approach is the Hardy–Littlewood maximal operator M defined by

$$Mf(x) = \sup_{r>0} \frac{1}{2r} \int_{-r}^r |f(x-t)| dt.$$

This, and thus the composition $M^k \equiv M \circ \dots \circ M$, is well-known to be bounded from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ with $p > 1$ (see for example the first theorem in [22]). The following lemma was proven with $M^3 w$ replaced by $(M|w|^q)^{1/q}$, with $q > 1$, in [9, 12] which was then precised in [24, 20]. This is sharp in the sense that the inequality does not necessarily hold if $M^3 w$ is replaced by $M^2 w$.

Lemma 2.1. *Let $\mathcal{T}$ denote the singular integral operator defined by*

$$\mathcal{T}f(x) = p.v. \int \mathcal{K}(x-y)f(y) dy,$$

where $\mathcal{K} \in C^1(\mathbb{R} \setminus \{0\})$ is such that $\mathcal{K}(x) \leq c_0|x|^{-1}$ and $|\frac{d}{dx}\mathcal{K}(x)| \leq c_0|x|^{-2}$. Then

$$\int |\mathcal{T}f|^2 w \leq C \int |f|^2 M^3 w,$$

where the constant C depends only on c_0 .

Ordering $\mathbb{S}^1$ in a clockwise sense, for each direction $\omega \in \Omega$ we write ω^{prev} for the previous direction of Ω and write $\omega^{\text{prev}} < \omega$. Considering the regions A_ω defined by

$$A_\omega = \left\{ \xi \in \mathbb{R}^3 : \omega^{\text{prev}} < \frac{(\xi_1, \xi_2)}{|(\xi_1, \xi_2)|} \leq \omega \right\},$$

we form the associated Fourier multiplier operators S_ω defined by

$$S_\omega f = \left(\chi_{A_\omega} \hat{f} \right)^\vee.$$

In the following lemma, we control the square function associated to these Fourier multipliers by two-dimensional directional maximal operators, for which we will need to recall some definitions.

The maximal operator M_ω is defined, initially on Schwartz functions, by

$$M_\omega f(x) = \sup_{r>0} \frac{1}{2r} \int_{-r}^r |f(x - t\omega)| dt.$$

This is bounded from $L^p(\mathbb{R}^3)$ to $L^p(\mathbb{R}^3)$, where $p > 1$, by an application of Fubini's theorem and the Hardy–Littlewood maximal theorem on the copy of the real line $\text{span}(\omega)$. For a set of directions $\Omega \subset \mathbb{S}^1 \subset \mathbb{R}^2$ we consider the directional maximal operator M_Ω defined by

$$M_\Omega : f \mapsto \sup_{\omega \in \Omega} M_\omega f,$$

and we also consider the strong maximal operator M_{st} defined by

$$M_{\text{st}}f(x) = \sup_{r_1, r_2 > 0} \frac{1}{4r_1 r_2} \int_{-r_1}^{r_1} \int_{-r_2}^{r_2} |f(x - t_1 e_1 - t_2 e_2)| dt_1 dt_2$$

which is bounded using Fubini's theorem as before.

We consider sequences $\{\theta_i\}_{i \in \mathbb{Z}}$ that satisfy $0 < \theta_{i+1} < \lambda \theta_i$ with lacunary constant $0 < \lambda < 1$. We say that a set of directions $\Omega \subset \mathbb{S}^1$ is lacunary (of order one) if there exists a lacunary sequence $\{\theta_i\}_{i \in \mathbb{Z}}$ so that there is at most one direction in each of the sets

$$\Omega_i = \left\{ \omega \in \Omega : \theta_{i+1} < \left| \frac{\omega_2}{\omega_1} \right| \leq \theta_i \right\}.$$

We are also obliged to consider the perpendicular directions

$$\Omega^\perp := \{ \omega \times e_3 \in \mathbb{S}^1 : \omega \in \Omega \},$$

and the associated maximal operator $M_{\Omega^\perp}$. Of course, by rotational symmetry, $M_{\Omega^\perp}$ is bounded if and only if M_Ω is bounded.

Lemma 2.2. *Let $\Omega \subset \mathbb{S}^1$ be lacunary. Then*

$$\int_{\mathbb{R}^3} \sum_{\omega \in \Omega} |S_\omega f|^2 w \leq C \int_{\mathbb{R}^3} |f|^2 M_{\text{st}}^3 M_{\Omega^\perp}^3 M_{\text{st}}^3 w,$$

where C depends only on the lacunary constant λ . Moreover,

$$\int_{\mathbb{R}^3} |f|^2 w \leq C \int_{\mathbb{R}^3} \sum_{\omega \in \Omega} |S_\omega f|^2 M_{\text{st}}^3 M_{\Omega^\perp}^3 M_{\text{st}}^3 w.$$

Proof. Let $\phi = \phi_o * \phi_o$ be a Schwartz function, supported on $[-\lambda^{-1/8}, -\lambda^{1/8}] \cup [\lambda^{1/8}, \lambda^{-1/8}]$, such that

$$(1) \quad \sum_{i \in \mathbb{Z}} \phi^2(\lambda^{i/8} \xi) = 1, \quad \xi \in \mathbb{R} \setminus \{0\},$$

where λ is the constant of lacunarity of the directions Ω . Consider the Fourier projection operators P_{ij} onto dyadic rectangles defined by

$$\widehat{P_{ij}f}(\xi) = \phi(\lambda^{i/8} \xi_1) \phi(\lambda^{j/8} \xi_2) \widehat{f}(\xi).$$

First we will prove that there is a constant C , depending only on λ , such that

$$(2) \quad \begin{aligned} \int_{\mathbb{R}^3} \sum_{i,j \in \mathbb{Z}} |P_{ij}f|^2 w &\leq C \int_{\mathbb{R}^3} |f|^2 M_{\text{st}}^3 w \\ \int_{\mathbb{R}^3} |f|^2 w &\leq C \int_{\mathbb{R}^3} \sum_{i,j \in \mathbb{Z}} |P_{ij}f|^2 M_{\text{st}}^3 w. \end{aligned}$$

This will follow by factorising into compositions of one-dimensional operators $P_{ij} = P_i^1 P_j^2$, in the obvious way, and applying Lemma 2.1.

Indeed, for all $t \in [0, 1]$, we first apply Lemma 2.1 to the operators

$$f \mapsto \sum_{i \in \mathbb{Z}} r_i(t) P_i^1 f,$$

where $r_i(t) = r_0(2^i t)$ and r_0 is the Rademacher function, equal to one on $[k, k + 1/2)$ and minus one on $[k - 1/2, k)$ for all $k \in \mathbb{Z}$. To see that the decay conditions of the lemma are satisfied, uniformly in $t \in [0, 1]$, we note that the associated kernel $\mathcal{K}$ satisfies

$$\mathcal{K}(x) = \sum_{i \in \mathbb{Z}} r_i(t) \lambda^{-i/8} \phi^\vee(\lambda^{-i/8} x) \approx \sum_{i \leq 8 \log_{\lambda-1} |x|^{-1}} r_i(t) \lambda^{-i/8} \phi^\vee(\lambda^{-i/8} x),$$

which follows by using the rapid decay of the Schwartz function $\phi^\vee$. We obtain that

$$\int_{\mathbb{R}^3} \left| \sum_{i \in \mathbb{Z}} r_i(t) P_i^1 g \right|^2 w \leq C \int_{\mathbb{R}^3} |g|^2 M_{e_1}^3 w,$$

where the constant is independent of $t \in [0, 1]$. By taking $g = \sum_j r_j(s) P_j^2 f$, in particular this implies that

$$\int_{\mathbb{R}^3} \left| \sum_{i, j \in \mathbb{Z}} r_i(t) r_j(s) P_{ij} f \right|^2 w \leq C \int_{\mathbb{R}^3} \left| \sum_{j \in \mathbb{Z}} r_j(s) P_j^2 f \right|^2 M_{e_1}^3 w,$$

where the constant C is independent of both $t \in [0, 1]$ and $s \in [0, 1]$. Applying Lemma 2.1 again, this time in the x_2 variable, yields

$$(3) \quad \int_{\mathbb{R}^3} \left| \sum_{i, j \in \mathbb{Z}} r_i(t) r_j(s) P_{ij} f \right|^2 w \leq C \int_{\mathbb{R}^3} |f|^2 M_{st}^3 w.$$

Integrating over $[0, 1] \times [0, 1]$ with respect to (t, s) yields the first inequality of (2) by the orthogonality of the Rademacher functions.

To see the reverse inequality, we take $f = \sum_{k, \ell} r_k(t) r_\ell(s) P_{k\ell} g$ and note that

$$\begin{aligned} & \sum_{i, j \in \mathbb{Z}} r_i(t) r_j(s) P_{ij} \left[\sum_{k, \ell \in \mathbb{Z}} r_k(t) r_\ell(s) P_{k\ell} g \right] \\ &= \sum_{i, j \in \mathbb{Z}} r_i^2(t) r_j^2(s) P_{ij} P_{ij} g + \sum_{i, j, k, \ell \in \mathbb{Z}: i \neq k, j \neq \ell} r_i(t) r_j(s) r_k(t) r_\ell(s) P_{ij} P_{k\ell} g \\ &= g + \sum_{i, j, k, \ell \in \mathbb{Z}: i \neq k, j \neq \ell} r_i(t) r_j(s) r_k(t) r_\ell(s) P_{ij} P_{k\ell} g, \end{aligned}$$

using (1) and that $r_i^2 r_j^2 \equiv 1$. Substituting into (3) and integrating over $[0, 1] \times [0, 1]$, we obtain

$$\int_{\mathbb{R}^3} \left(|g|^2 + \sum_{i, j \in \mathbb{Z}: i \neq j} |P_{ij} P_{ji} g|^2 \right) w \leq C \int_{\mathbb{R}^3} \sum_{k, \ell \in \mathbb{Z}} |P_{k\ell} g|^2 M_{st}^3 w.$$

by the orthogonality of the Rademacher functions as before. Note that unless $j = i \pm 1$, the sumands in the first sum are identically zero, but in any case they are positive and so this implies the second inequality of (2), for positive weights w . Then the inequality for positive weights implies the inequality for general weights.

Now our desired angular inequalities will follow by combining these more traditional Littlewood–Paley inequalities with a further application of Lemma 2.1 to the operator H_ω defined by

$$\widehat{H_\omega f}(\xi) = \chi_{\omega \cdot \xi \leq 0} \widehat{f}(\xi).$$




FIGURE 2. Segments with lacunary constant $1/2$ intersecting rectangles with lacunary constant $1/\sqrt{2}$.

Note that $H_\omega = \frac{1}{2}(\text{Id} + \frac{1}{i}\mathcal{H}_\omega)$, where $\mathcal{H}_\omega$ denotes the directional Hilbert transform defined by

$$\mathcal{H}_\omega f(x) = \frac{1}{\pi} p.v. \int f(x - \omega t) \frac{dt}{t}.$$

In the copy of the real line $\text{span}(\omega)$, this clearly satisfies the conditions of the lemma.

By enumerating the directions ω of Ω and writing r_ω for the associated Rademacher function we apply the inequalities of (2);

$$\begin{aligned}
 \int_{\mathbb{R}^3} \left| \sum_{\omega \in \Omega} r_\omega(t) S_\omega f \right|^2 w &\leq C \int_{\mathbb{R}^3} \sum_{i,j \in \mathbb{Z}} \left| \sum_{\omega: P_{ij} S_\omega \neq 0} r_\omega(t) S_\omega P_{ij} f \right|^2 M_{\text{st}}^3 w \\
 (4) \qquad &\leq C \sum_{i,j \in \mathbb{Z}} \int_{\mathbb{R}^3} |S_\omega P_{ij} f|^2 M_{\text{st}}^3 w \\
 &\leq C \int_{\mathbb{R}^3} \sum_{i,j \in \mathbb{Z}} |P_{ij} f|^2 M_{\Omega^\perp}^3 M_{\text{st}}^3 w \\
 &\leq C \int_{\mathbb{R}^3} |f|^2 M_{\text{st}}^3 M_{\Omega^\perp}^3 M_{\text{st}}^3 w,
 \end{aligned}$$

uniformly in $t \in [0, 1]$. Here, the second inequality is due to the fact that there are at most two directions ω which satisfy $P_{ij} S_\omega \neq 0$ for a given i and j (see Figure 2). The third inequality is by Lemma 2.1 in the directions $\pm\omega \times e_3$ or $\pm\omega^{\text{prev}} \times e_3$ as $S_\omega P_{ij}$ is equal to either $H_{\pm\omega \times e_3} P_{ij}$ or $H_{\pm\omega^{\text{prev}} \times e_3} P_{ij}$ (if it is not simply P_{ij}). Integrating over $[0, 1]$ with respect to t yields the first of our desired inequalities.

To see the reverse inequality, we take $f = \sum_{\omega'} r_{\omega'}(t) S_{\omega'} g$ as before, so that

$$\sum_{\omega \in \Omega} r_\omega(t) S_\omega \sum_{\omega' \in \Omega} r_{\omega'}(t) S_{\omega'} g = \sum_{\omega \in \Omega} r_\omega^2(t) S_\omega g = g,$$

as $r_\omega^2 \equiv 1$. Substituting into (4) we see that

$$\int_{\mathbb{R}^3} |g|^2 w \leq C \int_{\mathbb{R}^3} \left| \sum_{\omega' \in \Omega} r_{\omega'}(t) S_{\omega'} g \right|^2 M_{\text{st}}^3 M_{\Omega^\perp}^3 M_{\text{st}}^3 w,$$

and so by integrating over $[0, 1]$ with respect to t , we obtain the second inequality. $\square$

3. PROOF OF THEOREM 1.1

First of all, we give a precise version of the definition of finite order lacunary sets as mentioned in the introduction.

Definition 3.1. If Ω consists of a single direction we say that it is *lacunary of order zero*. We then say that $\Omega \subset \mathbb{S}^1 \subset \mathbb{R}^2$ is *lacunary of order K* if there is a choice of basis $\{e_1, e_2\}$ and lacunary sequence $\{\theta_i\}_{i \in \mathbb{Z}}$ such that the sets

$$\Omega_i = \left\{ \omega \in \Omega : \theta_{i+1} < \left| \frac{\omega_2}{\omega_1} \right| \leq \theta_i \right\}.$$

are lacunary of order $\leq K-1$ for all $i \in \mathbb{Z}$, with uniformly bounded lacunary constants.

Sjögren and Sjölin [21] proved that M_Ω is bounded from $L^p(\mathbb{R}^2)$ to $L^p(\mathbb{R}^2)$, where $p > 1$, if Ω is lacunary of finite order (see also [14, 18] for precedents and [2] for an equivalence).

Lemma 3.2. *If $\Omega \subset \mathbb{S}^1 \subset \mathbb{R}^2$ is lacunary of finite order and*

$$\Omega^{3d} := \left\{ \frac{1}{\sqrt{2}}(\omega \pm e_3) : \omega \in \Omega \right\},$$

then $M_{\Omega^{3d}}$ is bounded from $L^p(\mathbb{R}^3)$ to $L^p(\mathbb{R}^3)$ with $p > 1$.

Proof. We employ the localisation principle due to Parcet and the second author [19]. For directions $\Omega \subset \mathbb{S}^2 \subset \mathbb{R}^3$ and a lacunary sequence $\{\theta_i\}_{i \in \mathbb{Z}}$ we define segments Ω_i^ℓ by

$$\Omega_i^\ell = \left\{ \omega \in \Omega : \theta_{i+1} < \left| \frac{\omega_k}{\omega_j} \right| \leq \theta_i \right\}, \quad j < k,$$

where $e_\ell = e_j \times e_k$ or $-e_j \times e_k$. Then

$$(5) \quad \|M_\Omega\|_{p \rightarrow p} \leq C \sup_{\ell=1,2,3} \sup_{i \in \mathbb{Z}} \|M_{\Omega_i^\ell}\|_{p \rightarrow p},$$

where C depends only on $p > 1$ and the lacunary constant (see also [1] for a two-dimensional version). With this we can separate the directions of Ω^{3d} into isolated directions. The main difficulty is to choose the basis correctly between each application so that the lacunarity of Ω^{3d} is ‘visible’ in each direction.

Initially we change the basis vector e_3 by $e_3 \rightarrow \frac{1}{\sqrt{2}}(\omega_0 \pm e_3)$, where $\omega_0 \in \Omega \subset \mathbb{S}^1$ is one of the accumulation points of highest order, and take e_1 so that $e_3 + e_1$ is tangent to the circle which contains Ω^{3d} . This also fixes the final basis vector e_2 , chosen so that $e_3 = e_1 \times e_2$. Now, if Ω is lacunary of order K with lacunary constant λ , we can take $\theta_i = \lambda^{i/2}$ for all $i \in \mathbb{Z}$. Then the directions of Ω which correspond to the directions of the resulting segments of the localisation principle are lacunary of order $\leq K - 1$ (at least after eight applications of (5)), with lacunary constants uniformly bounded by λ . Thus, after a finite number of applications of (5), each time changing the basis as before to include one of the accumulation points of highest order, we separate the directions into isolated directions, and so $M_{\Omega^{3d}}$ is bounded. $\square$

Now as T_Ω is self-adjoint, the boundedness in L^p follows from the boundedness in $L^{p'}$, where $\frac{1}{p} + \frac{1}{p'} = 1$. To see that T_Ω is bounded in L^p with $2 < p < \infty$, by Hölder's inequality and the following weighted inequality it suffices to bound the maximal operators from $L^{(p/2)'}$ to $L^{(p/2)'}$. Thus, Theorem 1.1 is obtained by combining the following theorem with the previous lemma and the boundedness of $M_{\Omega \cup \Omega^\perp}$ due to Sjögren–Sjölin [21].

Theorem 3.3. *Let $\Omega \subset \mathbb{S}^1$ be lacunary of order K . Then*

$$\int_{\mathbb{R}^3} |T_\Omega f|^2 w \leq C \int_{\mathbb{R}^3} |f|^2 M_{\Omega \cup \Omega^\perp}^{15K} M_{\Omega^{3d}}^6 M_{\Omega \cup \Omega^\perp}^{15K} w,$$

where C depends only on the lacunary constant λ and the lacunary order K .

Proof. Noting that $M_\Omega = M_{\overline{\Omega}}$ we see that the relevant strong maximal operators (defined with respect to ω_0 and $\omega_0^\perp$, where ω_0 is an accumulation point) are dominated by $M_{\Omega \cap \Omega^\perp}^2$. Thus, by K applications of the second inequality of Lemma 2.2, we see that

$$\int_{\mathbb{R}^3} |T_\Omega f|^2 w \leq C \int_{\mathbb{R}^3} \sum_{\omega \in \Omega} |S_\omega T_\Omega f|^2 M_{\Omega \cup \Omega^\perp}^{15K} w.$$

Now we can write

$$S_\omega T_\Omega f = H_{\frac{1}{\sqrt{2}}(\omega - e_3)} H_{\frac{1}{\sqrt{2}}(\omega^{\text{prev}} - e_3)} S_\omega f + H_{\frac{1}{\sqrt{2}}(\omega + e_3)} H_{\frac{1}{\sqrt{2}}(\omega^{\text{prev}} + e_3)} S_\omega f,$$

so that, by two more applications of Lemma 2.1 in one of these four directions, we see that

$$\begin{aligned} \int_{\mathbb{R}^3} |T_\Omega f|^2 w &\leq C \sum_{\omega \in \Omega} \int_{\mathbb{R}^3} |S_\omega f|^2 M_{\Omega^{3d}}^6 M_{\Omega \cup \Omega^\perp}^{15K} w \\ &\leq C \int_{\mathbb{R}^3} |f|^2 M_{\Omega \cup \Omega^\perp}^{15K} M_{\Omega^{3d}}^6 M_{\Omega \cup \Omega^\perp}^{15K} w, \end{aligned}$$

where the second inequality is by K applications of the first inequality of Lemma 2.2. This completes the proof. $\square$

Remark 3.4. If we consider the two-dimensional multiplier operator R_Ω associated to a polygon $\mathcal{P}_\Omega$, defined by

$$R_\Omega : f \mapsto \left(\chi_{\mathcal{P}_\Omega} \hat{f} \right)^\vee,$$

where $\Omega \subset \mathbb{S}^1$ is lacunary of order K , the above argument also yields the weighted inequality

$$\int_{\mathbb{R}^2} |R_\Omega f|^2 w \leq C \int_{\mathbb{R}^2} |f|^2 M_{\Omega \cup \Omega^\perp}^{30K+6} w,$$

where the constant C depends only on the lacunary constant λ and the lacunary order K .

Remark 3.5. In higher dimensions, we are able to consider polytopes associated to lacunary directions with an exact product structure. That is to say that the orthogonal projections of the directions onto the two-dimensional subspaces, formed by the span of two basis vectors, are lacunary. For example, if we consider the n -dimensional multiplier operator

$$R_\Omega : f \mapsto \left(\chi_{\mathcal{P}_\Omega} \hat{f} \right)^\vee,$$

where $\Omega \subset \mathbb{S}^{n-1}$ consists of normalised versions of the directions

$$\{(2^{k_1}, \dots, 2^{k_n})\}_{k_1, \dots, k_n \in \mathbb{Z}},$$

the above argument also yields the weighted inequality

$$\int_{\mathbb{R}^n} |R_\Omega f|^2 w \leq C \int_{\mathbb{R}^n} |f|^2 M_\Omega^{30(n-1)+3 \cdot 2^{n-1}} w,$$

where the constant C depends only on the dimension. For this we first use $n-1$ applications of the second inequality of Lemma 2.2 (with $\mathbb{R}^3$ replaced by $\mathbb{R}^n$ and the lacunary sectors, divided by powers of 2, taken in $\text{span}(e_j, e_{j+1})$ with $j = 1, \dots, n-1$). Note that in this case the relevant two-dimensional strong maximal operators are bounded by M_Ω^2 . This separates the multiplier in such a way that we are left to deal with the composition of 2^{n-1} multiplier operators of the type that can be dealt with by Lemma 2.1 applied in one of the directions of Ω . We then recompose the operator with $n-1$ applications of the second inequality of Lemma 2.2. This maximal operator was bounded by Carbery [6], and so the multiplier associated to these polytopes with infinite faces are bounded on $L^p(\mathbb{R}^n)$ with $1 < p < \infty$.

The authors thank Jon Bennett for bringing [20] to their attention.

REFERENCES

- [1] A. Alfonseca, Strong type inequalities and an almost-orthogonality principle for families of maximal operators along directions in $\mathbb{R}^2$, J. London Math. Soc. **67** (2003), no. 1, 208–218.

- [2] M. Bateman, *Takeya sets and directional maximal operators in the plane*, *Duke Math. J.* **147** (2009), no. 1, 55–77.
- [3] J. Bennett, A. Carbery, F. Soria and A. Vargas, *A Stein conjecture for the circle*, *Math. Ann.* **336** (2006), no. 3, 671–695.
- [4] J. Bennett and S. Harrison, *Weighted norm inequalities for oscillatory integrals with finite type phases on the line*, *Adv. Math.* **229** (2012), no. 4, 2159–2183.
- [5] A. Carbery, *A weighted inequality for the maximal Bochner-Riesz operator on $\mathbb{R}^2$* , *Trans. Amer. Math. Soc.*, **287** (1985), no. 2, 673–680.
- [6] A. Carbery, *Differentiation in lacunary directions and an extension of the Marcinkiewicz multiplier theorem*, *Ann. Inst. Fourier.* **38** (1988), 157–169.
- [7] A. Carbery and A. Seeger, *Weighted inequalities for Bochner-Riesz means in the plane*, *Q. J. Math.* **51** (2000), no. 2, 155–167.
- [8] M. Christ, *On almost everywhere convergence of Bochner-Riesz means in higher dimensions*, *Proc. Amer. Math. Soc.* **95** (1985), 16–20.
- [9] R.R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, *Studia Math.* **51** (1974), 241–250.
- [10] A. Córdoba, *The Takeya maximal function and the spherical summation multipliers*, *Amer. J. Math.* **99** (1977), 1–22.
- [11] A. Córdoba, *An integral inequality for the disc multiplier*, *Proc. Amer. Math. Soc.* **92** (1984), no. 3, 407–408.
- [12] A. Córdoba and C. Fefferman, *A weighted norm inequality for singular integrals*, *Studia Math.* **57** (1976), no. 1, 97–101.
- [13] A. Córdoba and R. Fefferman, *On the equivalence between the boundedness of certain classes of maximal and multiplier operators in Fourier analysis*, *Proc. Nat. Acad. Sci. U.S.A.* **74** (1977), no. 2, 423–425.
- [14] A. Córdoba and R. Fefferman, *On differentiation of integrals*, *Proc. Nat. Acad. Sci. U.S.A.* **74** (1977), no. 6, 2211–2213.
- [15] C. Fefferman, *The multiplier problem for the ball*, *Ann. of Math. (2)* **94** (1971), 330–336.
- [16] S. Lee, K.M. Rogers and A. Seeger, *Improved bounds for Stein’s square functions*, *Proc. London Math. Soc.* **104** (2012), no. 6, 1198–1234.
- [17] S. Lee and A. Vargas, *On the cone multiplier in $\mathbb{R}^3$* , *J. Funct. Anal.* **263** (2012), no. 4, 925–940.
- [18] A. Nagel, E.M. Stein and S. Wainger, *Differentiation in lacunary directions*, *Proc. Nat. Acad. Sci. U.S.A.* **75** (1978), no. 3, 1060–1062.
- [19] J. Parcet and K.M. Rogers, *Differentiation of integrals in higher dimensions*, *Proc. Nat. Acad. Sci. U.S.A.* **110** (2013), no. 13, 4941–4944.
- [20] C. Pérez, *Weighted norm inequalities for singular integral operators*, *J. London Math. Soc. (2)* **49** (1994), no. 2, 296–308.
- [21] P. Sjögren and P. Sjölin, *Littlewood–Paley decompositions and Fourier multipliers with singularities on certain sets*, *Ann. Inst. Fourier* **31** (1981), 157–175.
- [22] E.M. Stein, *Singular Integrals and differentiability properties of functions*, Princeton University Press, Princeton, New Jersey, 1970.
- [23] E.M. Stein, *Some problems in harmonic analysis*, in *Harmonic analysis in Euclidean spaces* (Proc. Sympos. Pure Math., Williams Coll., Williamstown, Mass., 1978), Part 1, 3–20, Amer. Math. Soc., Providence, R.I.
- [24] J.M. Wilson, *Weighted norm inequalities for the continuous square functions*, *Trans. Amer. Math. Soc.* **314** (1989) 661–692.

DEPARTAMENTO DE MATEMÁTICAS – UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID,
SPAIN

E-mail address: `antonio.cordoba@uam.es`

INSTITUTO DE CIENCIAS MATEMÁTICAS CSIC-UAM-UC3M-UCM, 28049 MADRID, SPAIN

E-mail address: `keith.rogers@icmat.es`